Complete Summary

TITLE

Bipolar disorder: percentage of patients with Bipolar I Disorder with depressive symptoms and behaviors who have evidence of use of a mood stabilizing or antimanic agent during the first 12 weeks of pharmacotherapy treatment.

SOURCE(S)

STABLE (STAndards for BipoLar Excellence) performance measures. Boston (MA): Center for Quality Assessment and Improvement in Mental Health; 2007. various p.

Measure Domain

PRIMARY MEASURE DOMAIN

Process

The validity of measures depends on how they are built. By examining the key building blocks of a measure, you can assess its validity for your purpose. For more information, visit the Measure Validity page.

SECONDARY MEASURE DOMAIN

Does not apply to this measure

Brief Abstract

DESCRIPTION

This measure is used to assess the percentage of patients with Bipolar I Disorder with depressive symptoms and behaviors who have evidence of use of a mood stabilizing or antimanic agent during the first 12 weeks of pharmacotherapy treatment.

RATIONALE

Goals of Acute Treatment: Bipolar Disorder Depressive Episodes

- Remission of symptoms with a full return to usual levels of psychosocial functioning.
- Avoid mood destabilization and precipitating a manic or hypomanic episode.

Initial Treatment Considerations

- Treatment selection is dependent on illness severity, associated clinical features and patient preference.
- Treatment choice should also consider patient history, potential side effects and individual therapeutic response to specific pharmacotherapy agents.

Initial Treatment Recommendations

- Guidelines suggest lithium or lamotrigine alone or for patients refractory to monotherapy in combination with an atypical antipsychotic or lithium or lamotrigine alone or in combination, simultaneously, with an antidepressant.
- For patients refractory to the combinations above, guideline recommendations include the use of additional antimanic and antipsychotic agents.
- The use of antidepressants even in combination with a mood stabilizing or antimanic agent has been associated with some controversy over their potential to induce a mood switch.

PRIMARY CLINICAL COMPONENT

Bipolar I Disorder; depression; mood stabilizing agent; antimanic agent

DENOMINATOR DESCRIPTION

Patients with Bipolar I Disorder with symptoms or episodes that involve depression (see the related "Denominator Inclusions/Exclusions" field in the Complete Summary)

NUMERATOR DESCRIPTION

Patients with evidence of use of a mood stabilizing or antimanic agent during the first 12 weeks of pharmacotherapy treatment (see the related "Numerator Inclusions/Exclusions" field in the Complete Summary)

Evidence Supporting the Measure

EVIDENCE SUPPORTING THE CRITERION OF QUALITY

- A clinical practice guideline or other peer-reviewed synthesis of the clinical evidence
- A formal consensus procedure involving experts in relevant clinical, methodological, and organizational sciences
- One or more research studies published in a National Library of Medicine (NLM) indexed, peer-reviewed journal

Evidence Supporting Need for the Measure

NEED FOR THE MEASURE

Unspecified

State of Use of the Measure

STATE OF USE

Current routine use

CURRENT USE

Internal quality improvement

Application of Measure in its Current Use

CARE SETTING

Ambulatory Care Behavioral Health Care Physician Group Practices/Clinics

PROFESSIONALS RESPONSIBLE FOR HEALTH CARE

Advanced Practice Nurses
Physicians
Psychologists/Non-physician Behavioral Health Clinicians

LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED

Individual Clinicians

TARGET POPULATION AGE

Age greater than or equal to 18 years

TARGET POPULATION GENDER

Either male or female

STRATIFICATION BY VULNERABLE POPULATIONS

Unspecified

Characteristics of the Primary Clinical Component

INCIDENCE/PREVALENCE

Unspecified

ASSOCIATION WITH VULNERABLE POPULATIONS

Unspecified

BURDEN OF ILLNESS

See the "Rationale" field.

UTILIZATION

Unspecified

COSTS

Unspecified

Institute of Medicine National Healthcare Quality Report Categories

IOM CARE NEED

Getting Better Living with Illness

IOM DOMAIN

Effectiveness

Data Collection for the Measure

CASE FINDING

Users of care only

DESCRIPTION OF CASE FINDING

Patients with diagnosis of Bipolar I Disorder with depression: International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) or Diagnostic and Statistical Manual of Mental Disorders, Fourth Revision, Text Revision (DSM-IV-TR): 296.5x

DENOMINATOR SAMPLING FRAME

Patients associated with provider

DENOMINATOR INCLUSIONS/EXCLUSIONS

Inclusions

Patients 18 years of age or older with an initial diagnosis or new presentation/episode of bipolar disorder

AND

Documentation of Bipolar I Disorder with depressive symptoms, behaviors or episodes; to include at least one of the following:

- Code 296.5x documented in body of chart, such as a pre-printed form completed by a clinician and/or codes documented in chart notes/forms
- Diagnosis or impression documented in chart indicating Bipolar I Disorder with current depressive symptoms, behaviors or episodes

Exclusions

Exclude case from denominator population if numerator-required pharmacotherapy is not prescribed and the reason(s) for not prescribing is documented in chart, such as, not clinically indicated; not prescribed for related medical reasons (allergy, drug interaction); or not prescribed for patient reasons (patient refusal), etc.

RELATIONSHIP OF DENOMINATOR TO NUMERATOR

All cases in the denominator are equally eligible to appear in the numerator

DENOMINATOR (INDEX) EVENT

Clinical Condition Encounter

DENOMINATOR TIME WINDOW

Time window is a single point in time

NUMERATOR INCLUSIONS/EXCLUSIONS

Inclusions

Documented treatment with at least one pharmacotherapy agent with mood stabilizing or antimanic properties (Refer to the "Data Dictionary Reference" in the original measure documentation for specified medications.)

AND

Timeframe:

Documentation of the stated pharmacotherapy must be within the first 12 weeks of treatment.

Exclusions

Unspecified

MEASURE RESULTS UNDER CONTROL OF HEALTH CARE PROFESSIONALS, ORGANIZATIONS AND/OR POLICYMAKERS

The measure results are somewhat or substantially under the control of the health care professionals, organizations and/or policymakers to whom the measure applies.

NUMERATOR TIME WINDOW

Fixed time period

DATA SOURCE

Administrative data Medical record

LEVEL OF DETERMINATION OF QUALITY

Individual Case

PRE-EXISTING INSTRUMENT USED

Unspecified

Computation of the Measure

SCORING

Rate

INTERPRETATION OF SCORE

Better quality is associated with a higher score

ALLOWANCE FOR PATIENT FACTORS

Unspecified

STANDARD OF COMPARISON

Internal time comparison

Evaluation of Measure Properties

EXTENT OF MEASURE TESTING

- The STABLE measures were developed using the RAND Appropriateness Method and have been shown to have content validity and face validity.
- Data feasibility testing was performed to determine the availability of the data elements required in the measure numerator and denominator specifications.
- Inter-abstractor reliability testing was performed to assess the data collection strategy. The data collection strategy included data collection forms; data dictionary references and abstractor instructions.
- A field study was conducted to determine measure conformance in an appropriate convenience sample.

Refer to the references listed below for further information.

EVIDENCE FOR RELIABILITY/VALIDITY TESTING

STABLE performance measures: data feasibility testing & results. Boston (MA): Center for Quality Assessment and Improvement in Mental Health; 2007. 2 p.

STABLE performance measures: development process & validity ratings. Boston (MA): Center for Quality Assessment and Improvement in Mental Health; 2007. 3 p.

STABLE performance measures: field study process & conformance findings. Boston (MA): Center for Quality Assessment and Improvement in Mental Health; 2007. 3 p.

STABLE performance measures: inter-abstractor reliability testing & results. Boston (MA): Center for Quality Assessment and Improvement in Mental Health; 2007. 2 p.

Identifying Information

ORIGINAL TITLE

Bipolar disorder: use of mood stabilizing or antimanic agent in BD I with depression.

MEASURE COLLECTION

Standards for Bipolar Excellence (STABLE) Performance Measures

SUBMITTER

Center for Quality Assessment and Improvement in Mental Health

DEVELOPER

STABLE Project National Coordinating Council

FUNDING SOURCE(S)

AstraZeneca LLP, Wilmington, Delaware, provided financial sponsorship for the STABLE Project. They did not otherwise participate in the development of either the measures or toolkit.

COMPOSITION OF THE GROUP THAT DEVELOPED THE MEASURE

The STABLE <u>National Coordinating Council (NCC)</u> was comprised of national experts in bipolar disorder, psychiatry, primary care, and performance improvement. The NCC guided and directed the STABLE Project. NCC members agreed to serve with the understanding that the STABLE Performance Measures

and Resource Toolkit would be fully transparent and available without cost in the public domain.

<u>EPI-Q, Inc.</u>, is a consulting company providing practice-based outcomes research, pharmacoeconomic studies, and quality improvement services. EPI-Q managed the STABLE Project.

FINANCIAL DISCLOSURES/OTHER POTENTIAL CONFLICTS OF INTEREST

Unspecified

ADAPTATION

Measure was not adapted from another source.

RELEASE DATE

2007 Jan

MEASURE STATUS

This is the current release of the measure.

SOURCE(S)

STABLE (STAndards for BipoLar Excellence) performance measures. Boston (MA): Center for Quality Assessment and Improvement in Mental Health; 2007. various p.

MEASURE AVAILABILITY

The individual measure, "Bipolar Disorder: Use of Mood Stabilizing or Antimanic Agent in BD I with Depression," is published in "STABLE (STANdards for BipoLar Excellence) Performance Measures." This document is available in Portable Document Format (PDF) from the Center for Quality Assessment and Improvement in Mental Health (CQAIMH) Web site.

COMPANION DOCUMENTS

The following is available:

STABLE National Coordinating Council Resource Toolkit Workgroup. STABLE
resource toolkit. Boston (MA): Center for Quality Assessment and
Improvement in Mental Health; 2007 Mar. 67 p. This document is available in
Portable Document Format (PDF) from the Center for Quality Assessment and
Improvement in Mental Health (CQAIMH) Web site.

NQMC STATUS

This NQMC summary was completed by ECRI Institute on January 10, 2008. The information was verified by the measure developer on April 14, 2008.

COPYRIGHT STATEMENT

The STABLE materials published on this Web site are copyrighted by CQAIMH. They may be used for research, teaching, and quality measurement/improvement activities – provided the following:

- The materials are not sold, distributed or licensed for commercial purposes.
- CQAIMH's copyright is acknowledged in reproductions of these materials.
- Modifications to the materials are not made without CQAIMH's permission.

Disclaimer

NQMC DISCLAIMER

The National Quality Measures Clearinghouse™ (NQMC) does not develop, produce, approve, or endorse the measures represented on this site.

All measures summarized by NQMC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public and private organizations, other government agencies, health care organizations or plans, individuals, and similar entities.

Measures represented on the NQMC Web site are submitted by measure developers, and are screened solely to determine that they meet the NQMC Inclusion Criteria which may be found at

http://www.qualitymeasures.ahrq.gov/about/inclusion.aspx.

NQMC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or its reliability and/or validity of the quality measures and related materials represented on this site. The inclusion or hosting of measures in NQMC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding measure content are directed to contact the measure developer.

© 2008 National Quality Measures Clearinghouse

Date Modified: 11/3/2008

